



Vegetable Pro®
Herbicide
EPA Reg. No. 66222-15
For Control of Weeds on Direct Seeded
and Transplanted Celery
FIFRA 24(c) REGISTRATION
EPA SLN No. AZ-070003

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ARIZONA

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on this EPA-registered label are to be followed.
- This labeling must be in the possession of the user at the time of application.

For Agricultural or Commercial Use Only.

This SLN label is valid until December 31, 2016 or until otherwise amended, withdrawn, cancelled, or suspended.

DIRECTIONS FOR USE

CROP	TARGET PESTS	APPLICA-TION RATE	RATE AND METHOD OF APPLICATION	FREQUENCY OF APPLICATION
Celery (REI-24 hours)	Annual grass and broadleaf weeds	1.6 to 4.0 pints per acre	Apply in a minimum of 20 gallons of water per acre. Ground Application only.	<p>Direct-Seeded Celery: Within the rate ranges given, use the lower rates on coarse-textured soils and soils low in organic matter; use the higher rates on fine-textured soils and high in organic matter.</p> <ul style="list-style-type: none"> • Preemergence: Broadcast 2.4 to 3.2 pints per acre at planting or shortly after planting before celery emerges. • Postemergence: Broadcast 1.6 to 2.0 pints per acre after celery has 2-5 true leaves. Application may be made over the celery. Apply before weeds are 2 inches tall. <p>Transplants: Apply one application during the 2 to 6 week period after transplanting. Within the rate ranges given, use the lower rate on relatively coarse-textured soils and soils low in organic matter; use the higher rate on relatively fine-textured soils and soils high in organic matter. Application may be made over the celery. Apply before weeds are 2 inches tall.</p> <ul style="list-style-type: none"> • Coarse-textured soils: 2.0 to 3.2 pints per acre • Fine-textured soils: 3.2 to 4.0 pints per acre

OTHER REQUIREMENTS:

1. Do not apply by air or through any type of irrigation system.
2. **For Direct-Seeded Celery:** a) Make either one preemergence or one postemergence application (not both) per celery crop. b) Do not use on sand or loamy sand. c) Do not apply if celery is under water stress. d) Do not apply postemergence treatments of Vegetable Pro with other pesticides. Apply only after foliar applications of other pesticides are dry. e) Do not apply within two weeks after application of a herbicidal oil, such as "carrot" oil.
3. **Rotational Crops: Only the following crops** may be seeded 5 months after applying no more than 4 pints Vegetable Pro per acre on celery: cabbage, celery, corn, cotton, and peas. Onions and red beets may not be planted within 8 months of applying Vegetable Pro.
4. Do not enter treated area without protective clothing for 24 hours.
5. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas.

For fire, spill, and/or leak emergencies, contact Infotrac: 1-800-535-5053
For medical emergencies and health and safety inquiries, contact Prosar: 1-877-250-9291

24(c) REGISTRANT

Makhteshim Agan of North America, Inc.
3120 Highwoods Blvd, Suite 100
Raleigh, NC 27604

Vegetable Pro is a registered TM of Makhteshim Agan of North America, Inc. 2003

3/19/2012