



DEC 21 2012

INTERA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ARIZONA IN THE FOLLOWING COUNTIES – COCHISE, GRAHAM, LA PAZ, MARICOPA, MOJAVE, PIMA, PINAL, YUMA

FUSILADE® DX Herbicide

**EPA REG. No. 100-1070
EPA SLN No. AZ-070006**

For Control Of Annual And Perennial Grasses In Guayule: Non-food Use

This label is valid until December 31, 2017 or until otherwise amended, withdrawn, cancelled or suspended

KEEP OUT OF REACH OF CHILDREN

CAUTION

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Fusilade DX Herbicide label.

This label must be in the possession of the user at the time of application. The user must refer to the federal approved Fusilade DX label and read and follow direction for use, restrictions, and precautions.

WEEDS CONTROLLED

FUSILADE DX Herbicide may be applied as a post-emergence herbicide for control of annual and perennial grass weeds in guayule. Weeds controlled include the following.

- | | |
|-------------------------------------|---|
| Barnyard grass | Bermuda grass |
| Johnson grass, seedling and Rhizome | Volunteer cereals (barley, milo, oats, wheat) |

Use Directions		
Use Pattern	Rate Per Acre	Minimum Spray Volume Per Acre
Broadcast Application ¹	16-24 fl oz	Air or Ground 5 gallons

¹ Retreatment at the recommended rate may be needed to control late germinating grasses or if regrowth occurs.

APPLICATION DIRECTIONS

Ground/Aerial - Apply as a broadcast application in 5-40 gallons of water per acre by ground or 5-10 gallons of water per acre by air.

Spray Adjuvants - Always add one of the following spray adjuvants:

- **Crop oil concentrate** - Add a non-phytotoxic crop oil concentrate or a once refined vegetable oil containing 15-20% approved emulsifier at 0.5-1% v/v (1-2 pints per 25 gallons) of the finished spray volume. For aerial applications, add 1 pint of crop oil concentrate per acre.
- **Nonionic Surfactant** - Add nonionic surfactant containing at least 75% surfactant-active agent, at 0.25-0.5% v/v (1/2 -1 pint per 25 gallons) of the finished spray volume. For aerial applications, add 1 pint of surfactant per acre.

Refer to the EPA-registered label for additional information on application use directions.

SPECIFIC USE RESTRICTIONS

- DO NOT allow livestock to graze treated fields. DO NOT use treated crops for livestock feed. Crop residues are not to be fed to livestock or baled for such use. This information must be passed along to anyone using this crop.
- **DO NOT** apply a total of more than 72 fl. oz. of FUSILADE DX Herbicide per acre per 12 month period.

GENERAL RECOMMENDATIONS

- Refer to the approved EPA Sec. 3 label (Region B) for stage of weed growth and FUSILADE DX rate for best control.
- For best results, apply FUSILADE DX Herbicide 3 days before to 7 days after irrigation. In areas where tillage (discing, plowing, etc.) can not be performed, complete control of perennial grasses may not be possible.
- Make a second application to bermudagrass if re-growth occurs; usually about 4 weeks after first application. Bermudagrass control may be improved by directing the spray beneath the crop canopy. To improve coverage, make applications in a minimum of 20 gallons of water per acre.
- Make first application before the boot stage of Johnsongrass. If new shoots emerge or re-growth occurs, make a second application; usually about 4 weeks after first application.

24(c) registrant:
Syngenta Crop Protection, LLC.
P.O. Box 18300
Greensboro, NC 27419-8300

EPA SLN No.: AZ-070006

Label Code: AZ1070023BA1212