

## Section 24(c) Special Local Need Label

### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ARIZONA

GROUP 21 INSECTICIDE



#### EPA Reg. No 81880-4-10163 / EPA SLN NO. AZ-070009

#### For Control of Mites and Whiteflies on Greenhouse Tomatoes

ACTIVE INGREDIENT:	% By Wt.
PYRIDABEN: [2-tert-butyl-5-(4-tert-butylbenzylthio)-4-chloropyridazin-3(2H)-one]	75.0%
OTHER INGREDIENTS:	<u>25.0%</u>
	Total 100 0%

This label valid until December 31, 2017 or until otherwise amended, withdrawn, canceled or suspended

# WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

#### **DIRECTIONS FOR USE**

- It is a violation of Federal law to use this labeling in a manner inconsistent with its labeling.
- All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.
- This labeling must be in the possession of the user at the time of the pesticide application.

#### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Keep product and effluent water out of lakes, ponds, or streams with the exception of greenhouse wastewater holding tanks. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Do not apply when weather conditions favor drift from target area. Drift or runoff from treated areas may be hazardous to fish in adjacent sites. This product is highly toxic to bees. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Application early in the morning or at dusk is suggested.

CROP	RATE	PEST	COMMENTS		
Greenhouse	4 oz per 100 gallons of water	European red mite, Citrus red mite,	Apply when mites first appear and before a		
tomatoes	Or	Twospotted spider mite, Broad mite	threshold of five spider mites per leaf is reached.		
	0.09 oz per 1000 sq. ft.		·		
4–6 oz per 100 gallons of		Whiteflies			
	water				
0.09 - 0.14 oz per 1000 sq.					
	ft.				
	<ul> <li>Do not apply within 2 day of harvest (PHI)</li> <li>Do not make more than 2 applications per crop</li> <li>Do not apply more than 8 oz of product per crop</li> <li>Do not enter a treated greenhouse or a treated indoor area without protective equipment for 12 hours unless one of the following items is completed:         <ul> <li>10 air exchanges</li> <li>2 hours of system ventilation</li> <li>4 hours of ventilation using vents, windows or other passive ventilation</li> </ul> </li> </ul>				
	<ul> <li>All required</li> </ul>	d PPE is worn.			
	Allow a minimum of 30 days between sequential applications of <b>NEXTER</b> in crops that allow more than 1 application				
	per season.				
	<ul> <li>Do not apply this product t</li> </ul>	Do not apply this product through any type of irrigation system.			

Coverage: Apply Nexter in sufficient water to ensure thorough coverage of foliage and fruit. Thorough coverage is required for optimum control.

24(c) Registrant: Gowan Company

P.O. Box 5569 Yuma, AZ 85366-5569