

FIFRA 24(C) SPECIAL LOCAL NEED (SLN LABEL)

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF ARIZONA

ET[®] Herbicide/Defoliant

For Use on Guayule

**This label is valid until December 31, 2017
or until otherwise amended, withdrawn, canceled or suspended.**

Active Ingredient:

Pyraflufen ethyl: ethyl 2-chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1*H*-pyrazol-3-yl)-4-fluorophenoxyacetate.....**2.5%**

Other Ingredients*.....**97.5%**

Total:.....**100.0%**

Contains 0.208 lb. pyraflufen ethyl per gallon (25 grams per liter)

*contains petroleum distillates

EPA Reg. No. : 71711-7

EPA SLN No. : AZ-070011

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

All applicable directions, restrictions, and precautions on the EPA registered label are to be followed. This labeling must be in the possession of the user at the time of pesticide application.

GENERAL INFORMATION

For best results, use ET herbicide for control of annual broadleaf weeds when weeds are less than 4" in height or rosettes are less than 3" in diameter. ET is a contact herbicide. Thorough uniform spray coverage is essential for good control. Phytotoxicity may occur as leaf speckling; however, no crop loss is expected. **Addition of a crop oil concentrate (COC) or nonionic surfactant is recommended for optimum control.** Oil base spreaders will improve weed control but may also increase crop injury. Use with caution. Follow manufacturer's recommended use rates. Read and follow all label directions for each tank mix product. Always use in accordance with the most restrictive of label precautions and limitations.

SPRAY DRIFT

Avoid spray drift to all other crops and nontarget areas. Do not apply when weather conditions may cause drift. Do not allow this product to drift onto nontarget areas. Drift may result in illegal residues or injury to adjacent crops and vegetation, in the form of leaf yellowing and defoliation. To avoid spray drift, **DO NOT** apply aerially when wind speed is greater than 10 mph or during periods of temperature inversions. Use of larger droplet size will also reduce spray drift. **AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.**

APPLICATION RATE CHART

Guayule			
Pest	Rate / Acre	Spray Volume	Use Directions and Restrictions
Annual Broadleaf Weeds	1.0 to 2.75 oz	Minimum 20 gallons by ground; 10 gallons by air	<ul style="list-style-type: none"> Do not make more than 3 applications or exceed 5.5 fl oz/A per year for this use. Allow a minimum of 30 days between applications. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. NO portion of the treated field, including seed, seed screenings, or stubble, may be used for human consumption. This information must be passed along to anyone using this crop. Do not apply this product through any type of irrigation system.

ROTATIONAL CROP RESTRICTIONS

Crop/Crop Group	Rotational/Plantback Intervals
Corn Cotton Grapes Olives Pome Fruit Crop Group 11 Pomegranates Potatoes Soybeans Stone Fruit Crop Group 12 Tree Nuts Crop Group 14 Wheat, Triticale	0 days following application
Bulb Vegetables Crop Group 3 Cereal Grains Crop Group 15 (except corn, wheat, and triticale – see 0-day plantback interval above) Cole Crops Crop Group 5 Cucurbits Crop Group 9 Fruiting Vegetables Crop Group 8 Leafy Vegetables Crop Group 4 Legumes Crop Group 6 Oil Seeds Crop Group 20 Root and Tuber Vegetables Crop Group 1 (except potatoes – see 0-day plantback interval above) Sugarcane	1 day following preplant burndown application
For all other rotational crops, do not plant for 30 days following the last application of ET Herbicide/Defoliant.	

©2012 Nichino America, Inc.
 ET[®] is a trademark of Nichino America, Inc.

24c REGISTRANT
NICHINO
AMERICA
 4550 New Linden Hill Road
 Suite 501
 Wilmington, DE 19808
 1-888-740-7700
www.nichino.net