



The Chemical Company



Special Local Need

FOR DISTRIBUTION AND USE ONLY IN ARIZONA

FOR USE IN GUAYULE PRODUCTION IN LA PAZ, MARICOPA, MOHAVE, PIMA, PINAL AND YUMA COUNTIES ONLY

EPA Reg. No. 241-418

EPA SLN No. AZ-080009

Observe and follow all applicable directions, restrictions, worker protection standards, precautionary statements, mixing and application instructions on the EPA-registered label for Prowl® H2O herbicide, EPA Reg. No. 241-418.

around the plant to reduce the potential for the herbicide coming into contact with the roots of the guayule plants.

Apply the spray directly to the ground beneath the shrubs. Contact with leaves, shoots or buds by the spray mixture may cause malformed plant tissues.

Use rates for Prowl H2O alone and in tank-mix combinations are given in the following table.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application.

Table with 2 columns: Broadcast Rate Per Acre of Prowl (Quarts per Acre), Short term control (2.1 qts), Long-term control (4.2 qts)

Apply Prowl H2O at 2.1 to 4.2 quarts per acre during the spring and/or fall to control weeds expected to emerge during this period. DO NOT exceed 6.3 quarts/treated acre per year.

General Information

Prowl H2O may be applied for pre-emergence weed control of most annual grasses and certain broadleaf weeds in guayule crop production.

Chemigation Applications

Prowl H2O may be applied through sprinkler irrigation systems or flooded basin or furrow irrigation systems. DO NOT apply through any other type of irrigation system. Refer to and follow all recommendations, special instructions, and precautions in the Chemigation section on the Prowl H2O main container label.

Prowl H2O controls weeds as they germinate but will not control established weeds. Emerged weeds must be destroyed with tillage prior to applications or use Prowl H2O in combination with a post-emergence herbicide registered for use in guayule to remove the vegetation.

Prowl H2O treatments are most effective in controlling weeds when adequate rainfall or irrigation is received after application.

Use Methods, Rates and Timings

Pre-emergence - After transplanting guayule, uniformly apply in 20 or more gallons per acre (broadcast basis) with ground equipment. Applications may be band or broadcast.

Uniform distribution of Prowl-treated irrigation water is the sole responsibility of the applicator and is required to avoid crop injury, lack of herbicide effectiveness or illegal pesticide residues in the crop.

Apply after transplanting the guayule making sure that no roots are exposed and that the soil is well sealed

Application of **Prowl® H₂O herbicide** through irrigation systems should be used as a *supplemental weed control practice*. The addition of **Prowl H₂O** through irrigation systems will help prevent weed escapes at the irrigation point when the application is made before weed seeds germinate.

Prowl H₂O Tank Mixes

Prowl H₂O may be applied in a tank mix with any herbicide(s) that is registered for use in guayule to extend the weed spectrum. When using tank mixtures with **Prowl H₂O** always read the companion product(s) label to determine the specific use rates by soil types, weed species and weed growth stage. In addition, follow all precautions and restrictions including state and local restrictions that may apply to specific products. Always follow the most restrictive label.

Restrictions and Limitations

DO NOT apply by air.

DO NOT apply more than 6.3 quarts per acre per year (a single growing season).

DO NOT feed forage or graze livestock in treated areas.

DO NOT rotate to other crops for 24 months following a **Prowl H₂O** application to guayule.

CONDITIONS OF SALE AND WARRANTY

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

To the extent consistent with applicable law, BASF makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty.

To the extent consistent with applicable law, Buyer's exclusive remedy and BASF's exclusive liability, whether in contract, tort, negligence, strict liability, or otherwise, shall be limited to repayment of the purchase price of the product.

To the extent consistent with applicable law, BASF and the Seller disclaim any liability for consequential, special or indirect damages resulting from the use or handling of this product.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

0408

Prowl is a registered trademark of BASF.

© 2008 BASF Corporation.
All rights reserved.

000241-00418.20080421e.NVA 2008-04-195-0149

24(c) Registrant:
BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


The Chemical Company