

---

**RESTRICTED USE PESTICIDE**

**Due to Acute Toxicity**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

---

**TIERRA**

**Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ARIZONA IN THE FOLLOWING COUNTIES – COCHISE, GRAHAM, LA PAZ, MARICOPA, MOJAVE, PIMA, PINAL, YUMA**

**GRAMOXONE® SL 2.0**

**For Use on Guayule**

**This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2017**

**EPA Reg. No. 100-1431**

**EPA SLN No. AZ-120005**

**KEEP OUT OF REACH OF CHILDREN**

**DANGER/PELIGRO**

**POISON**



**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DESICCATION, ILLEGAL RESIDUES AND/OR CROP INJURY.**

---

**Directions for Use**

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered Gramoxone SL 2.0 label.

This label and the EPA registered label must be in the possession of the user at the time of pesticide application.

---

**Specific Crop Use Directions**

---

**Guayule (Pre and Postemergence Directed Spray and Harvest Aid)**

<b>Crop</b>	<b>Use Pattern</b>	<b>Rate Per Treated Acre</b>	<b>Minimum Spray Volume gal/acre</b>	<b>Preharvest Interval (Days)</b>
Guayule	Pre and Postemergence - Directed Spray	2.0 - 4.0 pts	Ground: 20 gals.	See precautions below
Desiccation of Guayule to facilitate harvest.	Harvest-Aid	2.0 - 4.0 pts.	Ground: 20 gals. Air 10 gals	

**ALWAYS ADD ONE OF THE FOLLOWING:**

Add a nonionic surfactant containing 75% or more surface-active agent at 0.125% v/v (1 pint per 100 gallons), OR add a nonionic surfactant containing 50-74% surface-active agent at 0.25% v/v (2 pints per 100 gallons) of the finished spray volume for ground applications. For aerial applications, add a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons) of the finished spray volume

**Specific Use Restrictions**

DO NOT allow livestock to graze treated fields. DO NOT use treated crops for livestock feed. Crop residues are not to be fed to livestock or baled for such use. If land used for guayule production changes ownership or use pattern, measures must be taken to ensure no crop residues are used for livestock feed. This information must be passed along to anyone using this crop.

## **Precautions, Restrictions, and Comments**

- DO NOT harvest until at least 4 days after application.
- Maximum number of applications per 12 month period is 5.
- DO NOT apply this product in a manner that allows spray to drift from the application target site and/or cause harm to humans, animals, or other non-target sites.
- DO NOT apply by ground equipment within 25 ft. or by air within 75 ft. of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries; and commercial fish ponds.
- Make aerial or ground applications when the wind velocity favors on target product deposition, (approximately 3-10 mph). Do not apply when wind velocity exceeds 10 mph for ground applications or 7 mph for aerial applications.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. Boom length must not exceed 75% of wingspan or rotor diameter.
- For directed spray applications, do not allow spray to contact green stems or foliage as injury may result.
- Use flat fan nozzles for the most effective application.
- Follow recommended spray volumes listed. These are minimum volumes only, and spray volumes should be increased as necessary to obtain complete coverage of the target weed or plant without runoff from the foliage.
- DO NOT apply this product through any type of irrigation system.
- DO NOT enter treated field within 12 hours of application.
- The user must refer to the federal approved GRAMOXONE SL 2.0 label and read and follow direction for use, restrictions, and precautions

Gramoxone® trademark of a Syngenta Group Company

© 2012 Syngenta

Section 24(c) Registrant:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, North Carolina 27419

AZ1431040AA0912